


1 / 16

FIG. 1


MLV-based transduction using Cre/loxP system as previously described



2 / 16

FIG. 2


EIAV-based transduction Cre/loxP system



3 / 16

FIG. 3

MLV SIN vector approach, with EIAV components in blue



4 / 16

FIG. 4

MLV-based transduction with HRE 3' LTR using Cre/loxP system





FIG. 5


MLV-based transduction for SIN vector production using Cre/loxP system



6 / 16


FIG. 6

MLV SIN-vector based transduction system. This general approach can be used with EIAV, HIV or MLV genomes



7 / 16

FIG. 7



8 / 16

FIG. 8







FIG. 9



9 / 16

**FIG. 10a**

FACS analysis of EV1 packaging cells prior to transduction with Trap2 vector

**FIG. 10b**

FACS analysis of EV1 packaging cell line transduced with Trap2 at an MOI of 0.3. A 5% top-slice of the highest expressers was carried out

10 / 16

FIG. 11

Validation of the $\Delta\Delta C_t$ method for quantitation of GFP mRNA, relative to β -actin.

A titration of total RNA from EV1 clone A was used. The difference in C_t values between the two assays is shown on the y-axis. The magnitude of the gradient must be <0.1 for the method to be valid. The gradient is 0.077, so the method is suitable.




FIG. 12

Quantitation of GFP mRNA relative to control β -actin mRNA. EV2 TD cells are transduced with Trap2 at an MOI of 0.3 and are the calibrator sample with the ratio designated 1.0.

Comparison of GFP expression levels in recombinase cell lines





FIG. 13



A) Original GFP expression of the clone.

B) GFP expression 7 days after transfection with Cre recombinase (pBS185). Excision frequency is 64%

C) Recombined clone 4 was identified as being negative for GFP


12 / 16




EV1 A4 cre/pONY8Z



EV2 D4 cre/pONY8Z



EV1 A4 pONY8Z



EV2 D4 pONY8Z

FIG. 14

13 / 16




FIG. 15

14 / 16

FIG. 16

Alignment of leader and gag regions present in vectors pONY4Z, 8Z and ATG mutated 8Z vector. The later is referred to as pONY8ZA. The sequence aligned are from the *NarI* site in the leader to the *XbaI* site between the EIAV gag sequence and the CMV promoter. Sequences in the leader are shown in *italic* and a space is present upstream of the position of the gag ATG.

4Z	1 <i>cgcccgaacagggacctgagaggggcgcagaccctacctgttgaacctgg</i>
8Z	1 <i>cgcccgaacagggacctgagaggggcgcagaccctacctgttgaacctgg</i>
mutated 8Z	1 <i>cgcccgaacagggacctgagaggggcgcagaccctacctgttgaacctgg</i>
4Z	51 <i>ctgacgtaggatccccgggacagcagaggagaacttacagaagcttct</i>
8Z	51 <i>ctgacgtaggatccccgggacagcagaggagaacttacagaagcttct</i>
mutated 8Z	51 <i>ctgacgtaggatccccgggacagcagaggagaacttacagaagcttct</i>
4Z	101 <i>ggagggtgttcctggccagaacacaggaggacaggtaag.at-gggagaccc</i>
8Z	101 <i>ggagggtgttcctggccagaacacaggaggacaggtaag.attgggagaccc</i>
mutated 8Z	101 <i>ggagggtgttcctggccagaacacaggaggacaggtaag.attgggagaccc</i>
4Z	150 <i>tttgacat-ggagcaaggcgctcaagaagttagagaaggtgacggtacaa</i>
8Z	151 <i>tttgacattggagcaaggcgctcaagaagttagagaaggtgacggtacaa</i>
mutated 8Z	151 <i>tttgacattggagcaaggcgctcaagaagttagagaaggtgacggtacaa</i>
4Z	199 <i>gggtctcagaaattaactactggtaactgtaattgggcgctaagtctagt</i>
8Z	201 <i>gggtctcagaaattaactactggtaactgtaattgggcgctaagtctagt</i>
mutated 8Z	201 <i>gggtctcagaaattaactactggtaactgtaattgggcgctaagtctagt</i>
4Z	249 <i>agacttatttcatt-gataccaactttgtaaaagaaaaggactggcagctg</i>
8Z	251 <i>agacttatttcatt-gataccaactttgtaaaagaaaaggactggcagctg</i>
mutated 8Z	251 <i>agacttatttcattgataccaactttgtaaaagaaaaggactggcagctg</i>

15 / 16

4Z	298	agggat-gtcattccattgctggaagat-gtaactcagacgctgtcagga
8Z	300	agggat-gtcattccattgctggaagat-gtaactcagacgctgtcagga
mutated 8Z	301	agggattgtcattccattgctggaagattgtaactcagacgctgtcagga
4Z	346	caagaaagagaggccttgaaagaacat-ggtgggcaatttctgctgtaa
8Z	348	caagaaagagaggccttgaaagaacat-ggtgggcaatttctgctgtaa
mutated 8Z	351	caagaaagagaggccttgaaagaacattggtgggcaatttctgctgtaa
4Z	395	agat-gggcctccagattaataat-gtagtagat-ggaaaggcatcattc
8Z	397	agat-gggcctccagattaataat-gtagtagat-ggaaaggcatcattc
mutated 8Z	401	agattgggcctccagattaataattgtagtagattggaaaggcatcattc
4Z	442	cagctcctaagagcgaaatat-gaaaagaagactgctaataaaaagcagt
8Z	444	cagctcctaagagcgaaatat-gaaaagaagactgctaataaaaagcagt
mutated 8Z	451	cagctcctaagagcgaaatattgaaaagaagactgctaataaaaagcagt
4Z	491	ctgagccctctgaagaatatct
8Z	493	ctgagccctctgaagaatatct
mutated 8Z	501	ctgagccctctgaagaatatct

FIG. 16 CONT'D

FIG. 17

Schematic representation of the structure of pONY 8.3G +/- vector genome plasmids

